

ONCE-DAILY DOSING



Take on an empty stomach upon waking



Take with a sip of water (no more than 120 mL)



Wait at least 30 minutes before eating, drinking, or taking any other oral medication

- Keep pill in the blister card until you are ready to take it, in order to protect it from moisture and light1
- Do not split, crush, or chew the pill when taking, as RYBELSUS® should be swallowed whole¹









STARTING DOSE¹

Start with 3 mg once daily for 1 month. This gives your body a chance to get used to the medicine

MAINTENANCE DOSES¹

Increase dose to 7 mg once daily for at least 1 month Based needs may dincread dose

Based on individual needs, your doctor may decide to increase your dose to 14 mg

- Packaging is conveniently colour-coded by dosage strength to avoid confusion
- Packaging for 3 mg strength is accented with green, 7 mg is accented with red, and 14 mg is accented with blue

Reference: 1. RYBELSUS® [summary of product characteristics]. Bagsværd, Denmark: Novo Nordisk A/S; April 2020.

Please see the Package Leaflet for more information or contact your doctor or nurse with any questions about how to manage your type 2 diabetes or use RYBELSUS®.



RYBELSUS® is a registered trademark of Novo Nordisk A/S. © 2020 Novo Nordisk A/S, Novo Allé, DK-2880, Bagsværd, Denmark HQ20RYB00034 Approval date: April 2020







